

PROTEOMIC ANALYSIS OF THE FUNGUS *Verticillium albo-atrum*

Stanislav MANDELIC¹, Sebastjan RADIŠEK², Polona JAMNIK³, Branka JAVORNIK⁴

¹Biotechnical Faculty, Department of Agronomy, Chair of Genetics, Biotechnology and Plant Breeding, Ljubljana

²Slovenian Institute for Hop Research and Brewing, Žalec

³Biotechnical Faculty, Food Science and Technology Department, Chair of Biotechnology, Ljubljana

ABSTRACT

Verticillium wilt caused by the soilborne phytopathogenic fungus *Verticillium albo-atrum* is a serious threat to hop production in Slovenia. The fungus invades the plant through its roots and colonizes the vascular tissue, which leads to a disrupted or entirely blocked water supply. Mild and lethal pathotypes are distinguished, depending on the level of virulence. The mild form has been present in Slovenia since 1974 and causes no considerable economic loss, while the lethal form was first discovered in 1997. Due to the lack of effective phytopharmaceutical agents, the only effective means of fighting the disease are phytosanitary and hygienic measures, crop rotation and planting resistant varieties. In order to get a wide insight into the infectious process and to identify proteins related to infection, we employed a proteomic approach, which allows simultaneous analysis of total proteins on two-dimensional polyacrylamide gel. The results of comparison of total proteins between the mild and lethal forms are presented.

Key words: 2D electrophoresis, hop, *Humulus lupulus*, *Verticillium albo-atrum*

IZVLEČEK

PROTEOMSKA ANALIZA GLIVE *Verticillium albo-atrum*

Hmeljeva uvelost, ki jo povzroča talna fitopatogena gliva *Verticillium albo-atrum*, predstavlja resno grožnjo pridelavi hmelja v Sloveniji. Gliva skozi korenine vdre v rastlino in se razmnožuje v prevajalnem tkivu, kar privede do motene ali povsem blokirane oskrbe z vodo. Glede na stopnjo virulentnosti ločimo blag in letalen patotip glive. Medtem ko je blaga oblika v Sloveniji zastopana od leta 1974 in ne povzroča večje gospodarske škode, pa je bila letalna oblika, ki rastline popolnoma uniči, odkrita leta 1997. Ker učinkovitih fitofarmaceutskih pripravkov za zatiranje glive in zdravljenje obolelih rastlin ni, so fitosanitarni in higienski ukrepi, prilagojen kolobar in gojenje odpornih sort edina učinkovita sredstva v boju proti bolezni. Da bi dobili širok vpogled v proces infekcije in identificirali proteine, ki pri tem sodelujejo, smo uporabili proteomski pristop, ki omogoča hkratno analizo celokupnih proteinov na poliakrilamidnem gelu v dveh dimenzijah. Rezultati primerjave celokupnih proteinov iz blage in letalne oblike bodo predstavljeni v nadaljevanju.

¹ Jamnikarjeva 101, SI-1111 Ljubljana

² dr. agr. znan., Cesta Žalskega tabora 2, SI-3310 Žalec

³ dr., Jamnikarjeva 101, SI-1111 Ljubljana

⁴ dr., prav tam

Ključne besede: 2D elektroforeza, hmelj, *Humulus lupulus*, *Verticillium albo-atrum*

1 INTRODUCTION

Verticillium wilt is a vascular disease caused by the soil borne pathogenic fungi *Verticillium albo-atrum* and *Verticillium dahliae*. *V. albo-atrum* isolates infecting hop display different levels of virulence, resulting in mild or lethal symptoms. Slovene isolates have been determined as PG1 (mild) or PG2 (lethal) pathotype based on pathogenicity tests and AFLP analysis. The spread of the lethal form in hop gardens in the Savinja Valley is a serious threat to hop production. From 1997 to 2006, more than 180 ha of hop gardens were infected and almost 90 ha were destroyed. Due to the lack of effective phytopharmaceutical agents, the only effective means of fighting the disease are phytosanitary and hygienic measures, crop rotation and planting resistant varieties. We employed a proteomic approach to gaining a wide insight into the infection process and to identifying proteins related to infection. This allows simultaneous analysis of total proteins on a two-dimensional polyacrylamide gel.

2 MATERIALS AND METHODS

Verticillium albo-atrum was grown in a general fungal medium for 1 week and mycelium was collected by filtration. Proteins were extracted according to Jamnik *et al.*, 2006, and the concentration was measured with a Bradford reagent. 450 µg of proteins were precipitated, resuspended in rehydration solution and applied to 13 cm IPG strips (pH 4-7) by rehydration loading. After isoelectric focusing, strips were equilibrated and loaded on 12.5% polyacrylamide gels. After electrophoresis, gels were stained with Coomassie Brilliant Blue G-250, as described by Neuhoff *et al.*, 1985, and scanned. Dymension 2 software was used for image analysis and statistical tests. Four biological and three technical replicates were included in the experiment (a total of 24 gels).

3 RESULTS

Separation of proteins was good and there were only a few spots with horizontal or vertical streaks. Reproducibility between technical replicates was excellent, while biological replicates displayed minor differences. On average, 2645 spots were detected in PG1 (660 valid) and 2529 spots in PG2 (721 valid). A spot was determined as valid if it was present on at least 9 replicate gels (out of 12) and if it passed filtering criteria. There were 173 spots present only in the PG1 pathotype and 234 spots were found only in the PG2 pathotype; 487 spots were observed in both pathotypes. A regulation factor of 2 was set, which resulted in 64 spots significantly more abundant in the PG1 pathotype and 90 spots significantly more abundant in the PG2 pathotype ($p < 0,05$). Twenty-seven spots were chosen for subsequent MS analysis (Fig. 1 and Fig. 2).




Fig. 1: Representative image of 2-D gel (PG1 pathotype). Spots chosen for MS analysis are marked by arrows.




Fig. 2: Representative image of 2-D gel (PG2 pathotype). Spots chosen for MS analysis are marked by arrows.

4 CONCLUSION

Two-dimensional electrophoresis was used successfully for the separation of cellular proteins from *Verticillium albo-atrum*. Significant differences between PG1 and PG2 pathotypes were observed and will be further investigated. Furthermore, the general fungal medium will be replaced by a simulated xylem fluid medium (SXM) to induce the expression of genes related directly to infection.

5 REFERENCES

- Radišek, S., Jakše, J., Simončič, A., Javornik, B., 2003. Characterization of *Verticillium albo-atrum* field isolates using pathogenicity data and AFLP analysis. *Plant Disease*, 87: 633-638
- Radišek, S., 2006. Hmeljeva uvelost v slovenskih hmeljiščih. Inštitut za hmeljarstvo in pivovarstvo Slovenije: 24 str.
- Jamnik, P., Radišek, S., Javornik, B., Raspor, P., 2006. 2-D separation of *Verticillium albo-atrum* proteins. *Acta agriculturae Slovenica*, 87-2: 455-460
- Neuhoff, V., Stamm, R., Eibl, H., 1985. Clear background and highly sensitive protein staining with Coomassie Blue dyes in polyacrylamide gels: A systematic analysis. *Electrophoresis*, 6: 427-448